

## Mantle-earthquake geothermometry of rejuvenated Proterozoic lithosphere, western Saudi Arabia

Alexander R. Blanchette<sup>1,2</sup>, Simon L. Klemperer<sup>1</sup>, Walter D. Mooney<sup>2</sup>, Hani M. Zahran<sup>3</sup>


<sup>1</sup>Department of Geophysics, Stanford University, Stanford, CA 94305-2215, U.S.A., [sklemp@stanford.edu](mailto:sklemp@stanford.edu)

<sup>2</sup>Earthquake Science Center, United States Geological Survey, Menlo Park, CA 94025, U.S.A.

<sup>3</sup>Saudi Geological Survey, Jeddah 21514, Saudi Arabia

Harrat Lunayyir is a Quaternary volcanic field above a 2009 dyke intrusion located in the Arabian Shield ~100 km east of the Red Sea rift. We use 3D common conversion point stacking of P-wave receiver functions to show that the Moho is at  $38 \pm 2$  km depth, close to the 40-km crustal thickness measured in the center of the Arabian craton, whereas the lithosphere-asthenosphere boundary is at  $60 \pm 5$  km depth (Figure 1), far shallower than the 150 km seen further east in the Arabian shield. We re-located 64 high-frequency earthquakes with  $m_L \leq 2.5$  at depths of 42–48 km, clearly beneath the Moho, ~30 km east-southeast of the 2009 dyke-intrusion beneath Harrat Lunayyir (Figure 1). These brittle-failure earthquakes, that must have nucleated at temperatures  $T \leq 900^\circ\text{C}$ , show that the lithosphere is not in thermal equilibrium with the underlying asthenosphere that has a petrologically calibrated potential temperature of  $\sim 1350^\circ\text{C}$  at 60 km depth (Blanchette et al., 2018).

We take the maximum depth of crustal earthquakes as marking the crustal brittle-ductile transition, thereby constraining the  $350 \pm 50^\circ\text{C}$  isotherm. We take the maximum depth of sub-Moho earthquakes as marking the upper-mantle ductile-brittle transition, thereby constraining the  $850 \pm 50^\circ\text{C}$  isotherm. We can then model the thermal evolution of the shield margin from a stable cratonic geotherm (derived from observed surface heat-flow) prior to onset of Red Sea extension, while the lithosphere is eroded from  $\geq 125$  km to its modern 60-km thickness. Absence of earthquakes below 48 km constrains the lithosphere to have been thinned to 60 km no earlier than  $\sim 12$  Ma, assuming earthquakes cannot nucleate at  $T \geq 900^\circ\text{C}$ . Absence of crustal earthquakes below 20 km constrains the lithosphere to have been thinned to 60 km no later than  $\sim 11$  Ma, otherwise the aseismic lower crust would encompass temperatures  $\leq 300^\circ\text{C}$ . Initial Red Sea extension at  $\sim 30$  Ma must have been followed by a second phase of lithospheric thinning continuing until  $\sim 12 \pm 2$  Ma, consistent with onset of Younger Harrat magmatism and focusing of extension towards the Red Sea rift axis from  $\sim 12$  Ma (Blanchette et al., 2018).



**Figure 1.** 3d CCP stack of P-wave receiver functions and re-located sub-Moho earthquakes (projected on to cube faces, within circles), with 2-km voxels. Blue (red) color indicates wave-speed decrease (increase) with depth. Vertical lines: locations of slices through the cube that are displayed on the west-east and south-north cube faces. Top face: geologic map. Surface trace of 2009 dyke: red line; Quaternary basalts: grey; Proterozoic basement: yellow. Graph: inferred modern non-equilibrium thermal gradient compared to pre-Neogene cratonic geotherm.

**References**

Blanchette, A.R., Klemperer, S.L., Mooney, W.D., and Zahran, H.M., 2018, Two-stage Red Sea rifting inferred from mantle earthquakes in Neoproterozoic lithosphere, *Earth and Planetary Science Letters*, 497, 92-101.