

Geodynamic and geophysical consequences of craton: Implications for metallogeny

Suzanne Y. O'Reilly¹, William L. Griffin¹

¹ARC Centre of Excellence for Core to Crust Fluid Systems (CCFS) and GEMOC, Earth and Planetary Sciences, Macquarie University, NSW 2109, Australia, sue.oreilly@mq.edu.au, bill.griffin@mq.edu.au

Introduction

Convergence between geochronology datasets of Hf isostopic model ages for zircons (e.g., Belousova et al., 2010) and Re-Os model ages for mantle sulfides (Griffin et al., 2014), reinforced by other geochemical and tectonic criteria, indicates that over 75% of the subcontinental lithospheric mantle (SCLM) and its overlying crust (now mostly lower crust) formed at ~2.9-3.5 Ga, probably in global overturn events that marked a change in Earth's fundamental geodynamic behaviour (Griffin et al., 2014; Griffin and O'Reilly, 2017). *This is a critical finding for understanding the location and type of metallogenic deposits associated with mantle-derived fluids and magmas.*

Pristine Archean lithospheric mantle, the roots of the Archean cratons up to >300km deep, is not only *more* depleted (low in basaltic melt components) than younger lithospheric mantle, it is *differently* depleted. Garnet-peridotite suite rocks in cratonic mantle are now interpreted as products of the metasomatic introduction of garnet and clinopyroxene into original depleted harzburgite, thus “refertilising” the depleted residue. *Therefore, the “oceanic melting trend” is now re-interpreted as a refertilisation trend, with the compositional trend arrow reversed* (Griffin et al., 2009). The cratonic roots in contact with the asthenosphere, and their vertical margins in contact with repeated fluid and melt fluxes (O'Reilly et al., 2009), are the most strongly modified domains through time, so cratonic root compositions show a general increase in fertility with increasing depth (Figure 1).

Integration of geochemical and geophysical datasets, groundtruthed with petrophysical measurements and modelling for different mineral compositions and modes, has enabled geologically realistic interpretations of global (and regional) tomographic results, identifying the vertical and horizontal extent of original Archean cratonic mantle and their refertilised domains (Griffin et al., 2009).

Stealth Metasomatism and its Geophysical Significance

Metasomatism (and associated refertilisation) affects not only the geochemistry of the lithospheric mantle, but also its physical *parameters* (and hence geophysical signatures): e.g. density, seismic response and thermal and electrical characteristics. The concept of “*stealth*” metasomatism has been introduced to highlight the “deceptive” addition to lithospheric mantle rock-types of new phases (e.g. garnet and/or clinopyroxene) indistinguishable mineralogically from common mantle-peridotite phase assemblages (O'Reilly and Griffin, 2012). Recognition of *stealth metasomatism* reflects the increasing awareness of the importance of refertilisation of ancient refractory mantle regions by metasomatic fluid fronts in modifying mantle domains. *Understanding the timing and nature of stealth metasomatism is critical to understanding the geochemical and geodynamic evolution of the lithospheric mantle domains and assessing their metallogenic fertility.*

Primitive Archean lithospheric mantle is highly magnesian (~49% MgO) with lower density (~3.31 g/cm³) than fertile mantle (~3.37 g/cm³). The resulting contrasts in *seismic response* become measurable if metasomatised regions are on the scale of tens to hundreds of km. Metasomatic refertilisation of the cratonic lithospheric mantle not only increases its density, but also strongly affects its rheology. *The recognition that the cratonic lithospheric mantle may in general consist of a very depleted upper layer, becoming more metasomatised (evidenced by garnet peridotites) at depth, provides a solution to some important discrepancies between geophysical data and numerical models.* A deep cratonic root made up mainly of garnet peridotites would imply a lower geoid and a much greater elevation than is observed for

the Kaapvaal craton, and a mismatch in Vp/Vs ratios (Afonso et al., 2010), but a layered model in with refertilization increasing with depth, yields numerical models that fit the geophysical data. *Thermal changes* are caused by advective transfer of heat by relatively hot metasomatising fluids and by the influx of heat-producing elements (K, U, Th) that accompanies some types of metasomatism. Radioactive decay of these elements can raise the local heat flow by 50-70% over normal reduced mantle heat flow. The origins of significantly contrasting *electromagnetic (MT) responses* in the lithospheric mantle have not yet been satisfactorily verified. Fluid-related recrystallisation demonstrated in microstructural studies may result in significant H contents in mantle minerals may affect electrical conductivity. Comparisons of high-resolution MT traverses with geochemical mapping traverses show a correlation of higher conductivity with more strongly metasomatised domains.

Implications for metallogeny

Magma-related ore systems form economic deposits that underpin our human civilisation. The magmas related to metallic element redistribution derive from the asthenosphere, then traverse and interact to varying degrees with the subcontinental lithospheric mantle. The evolution of the original Archean lithospheric mantle has been the single largest influence on the formation of most of Earth's ore deposits (Griffin et al., 2013):

- the high degree of buoyancy of this ancient SCLM relative to the asthenosphere, results in the persistence today of low-density, rheologically coherent Archean domains and commonly, the preservation of old crustal (or at least lower-crustal) domains;
- the enduring (and volumetrically dominating) Archean lithospheric mantle domains are a reservoir for metasomatic enrichment over their long history, creating a potentially metallogenically-fertile mantle impregnated with critical elements (e.g., Au, Cu, Ni and PGEs (Zhang et al., 2008);
- the formation of Archean cratons provided an architectural mantle-scape of regions with contrasting rheology, composition and thickness. These cohesive Archean domains direct magma and fluid pathways around their margins and along old sutures between blocks, acting as both sinks and sources for ore-forming elements depending on the geodynamic stage (Griffin et al., 2013)
- if the first stabilisation of lithospheric mantle at 3.0-3.5 Ga signalled the end of a mantle overturn regime (either uniquely, or intermittently with subduction), then this is when long-lived tectonic regimes conducive to mineralising systems (e.g. back-arc basins, passive margins, cratonic boundaries) became available (Griffin et al., 2013; Begg et al., 2017).




Figure 1. Interactions between magmas and the base and edges of cratonic roots. Plume magmatism triggers kimberlite formation and flows to areas of thinner SCLM where melting is focused. Variable interaction of melts with crust and SCLM influences Ni-Cu and PGE deposit genesis (Griffin et al., 2013). Lithosphere-scale discontinuities and zones of weakness can focus the locus of kimberlite eruption.

References

- Afonso J.C., Ranalli G., Fernandez M., et al. 2010, On the Vp/Vs-Mg# correlation in mantle peridotites: Implications for identification of thermal and compositional anomalies in the upper mantle, *Earth Planet Sci Lett*, 289, 606-618.
- Begg, G.C., Hronsky, J.M.A., Griffin, W.L. and O'Reilly, S.Y., 2017, Global- to District-Scale Controls on Orthomagmatic Ni-Cu-(PGE) and PGE Reef Deposits, Mondal, S. and Griffin W.L. (eds.) *Processes and ore deposits of ultramafic-mafic magmas through space and time*. Elsevier. pp. 1-46.
- Belousova E.A., Kostitsyn Y.A., Griffin W.L., Begg, G.C., O'Reilly, S.Y. et al., 2010, The growth of the continental crust: Constraints from zircon Hf-isotope data. *Lithos* 119, 457-466.
- Griffin, W., O'Reilly, S.Y., Afonso, J.C., Begg, G.C., 2009. The composition and evolution of lithospheric mantle: A re-evaluation and its tectonic implications. *J. Petrology* 50(9), 1185-1204.
- Griffin W.L. et al., 2014, The world turns over: Hadean-Archean crust-mantle evolution, *Lithos* 189:, 2-15.

- Griffin, W.L. and O'Reilly, S.Y. 2017. The Earliest Subcontinental Mantle. In M. Van Kranendonk, H. Smithies and V. Bennett (eds.) *Earth's Oldest Rocks*. 2nd edition. Elsevier *in press* Sept. 2018.
- Griffin W.L, Begg G.C., O'Reilly S.Y., 2013, Continental-root control on the genesis of magmatic ore deposits. *Nature Geoscience* 6: 905-910.
- O'Reilly, S.Y. and Griffin, W.L., 2012., Mantle Metasomatism, in Harlov, D.E. and Austrheim H. (eds.) *Metasomatism and the Chemical Transformation of Rock*, Springer-Verlag Berlin Heidelberg 2012. Pp 467-528.
- O'Reilly S.Y., Zhang M, Griffin W.L., Begg G., Hronsky J., 2009, Ultradeep continental roots and oceanic remnants: a solution to the geochemical "mantle reservoir" problem? *Lithos* 112(2), 1043-1054.